

Jay L. Cohen, M.D., P.C.

464 Hillside Avenue, Suite 303
Needham, MA 02494
Phone: (781) 449-3588
Fax: (781) 449-5474

2000 Washington Street, Suite 404
Newton, MA 02462
Phone: (781) 449-3588
Fax: (781) 449-5474

Facial Acne – Topical Antibiotic and Retinoid

You have been prescribed two topical medications for the treatment of facial acne. This sheet will help you understand how to use these medications and the best practices to follow. Please review this information before starting your medications.

In the **morning** apply _____, which is a **topical antibiotic**.

At **night** apply _____, which is a **retinoid** (a chemical related to Vitamin A. The original medication was known as Retin-A). The retinoid product may be inactivated by sunlight, so it is better to apply this at night.

If your schedule makes it difficult to apply a cream/lotion in the morning, you can still apply the medications twice a day. The “morning” medication can be applied in the late afternoon or early evening. A few hours later you can wash off the “morning” medication and the “night” medication can be applied. Similarly, if you do not like to sleep with a cream on your skin you may apply it a few hours before bedtime and then wash it off before you go to sleep. The key point is this: as long as each cream is on your skin for 1-2 hours, you will get the full benefit.

How to apply the medications:

- Before applying these medications, wash your face with a mild soap, pat dry, and allow it to dry completely. For the antibiotic, apply a thin layer evenly over your skin. For the retinoid, apply one pea-sized amount of cream to your fingertip, dab it on your forehead, nose, cheeks and chin, then smooth it into a thin even layer over your whole face (avoiding the eye area). The retinoid cream should disappear almost immediately as you massage it into your skin. You may require a little more or less of each medication, so adjust the amount you apply to your particular need.
- If it is helpful, you may add 1-2 pea-sized portions of moisturizer to the retinoid cream so that it will be easier to evenly cover the entire area needing treatment.

When you are using these medications for the first time, there are a few important steps to follow, and helpful hints to keep in mind:

- Do not apply both the topical antibiotic and the retinoid medications at the same time. Always apply them at different times, at least a few hours apart.

Jay L. Cohen, M.D., P.C.

464 Hillside Avenue, Suite 303
Needham, MA 02494
Phone: (781) 449-3588
Fax: (781) 449-5474

2000 Washington Street, Suite 404
Newton, MA 02462
Phone: (781) 449-3588
Fax: (781) 449-5474

- Do not use more than recommended or apply more frequently than instructed, because this may cause irritation, and will not provide faster or better results.
- Apply these medications anywhere you have acne or tend to get it, even if these areas have no acne at this time. These medications will do a reasonably good job of healing existing acne, but they do an even better job of preventing new acne.
- Your skin may be sensitive to these medications when you begin to use them. Side effects may include discomfort, peeling, and a slight redness of the skin. Redness or peeling usually subsides within two to four weeks. If necessary, the medication can be applied every other day until your skin can tolerate a daily application. Do not apply these medications to the corners of the nose, mouth, or eyes, or to any open sores.
- Please have patience. Your skin may get worse before it gets better. This is normal. Do not worry! When the medications are used properly, improvement is usually noted within four to six weeks, with considerable improvement noticeable in twelve to fourteen weeks.
- Do not stop using your medications once you begin to see improvement. Continued use is needed to prevent recurrence of your acne. Once your acne is under control, you should continue to use your medications until Dr. Cohen instructs you to do otherwise.

Side Effects

Initially you may have some peeling and redness of your skin, which is normal. If the peeling or redness is severe, please discontinue the medication and contact us at the phone number listed below.

Your skin may become more sensitive to the sun. Use a non-greasy (noncomedogenic) sunscreen, with a minimum of SPF 30, even when it is cloudy out. You should also avoid exposure to wind or excessive cold.

****It is recommended that you not use these medications if you may be pregnant or are planning to become pregnant. ****